

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
12 April 2001 (12.04.2001)


PCT

(10) International Publication Number  
WO 01/25454 A2

- (51) International Patent Classification<sup>7</sup>: C12N 15/82, 9/02 (74) Agents: CÔTÉ, France et al.; Swabey Ogilvy Renault, Suite 1600, 1981 McGill College Avenue, Montréal, Québec H3A 2Y3 (CA).
- (21) International Application Number: PCT/CA00/01143
- (22) International Filing Date: 2 October 2000 (02.10.2000) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/157,133 4 October 1999 (04.10.1999) US
- (71) Applicant (*for all designated States except US*): MED-ICAGO INC. [CA/CA]; 2480, rue Hochelaga, Sainte-Foy, Québec G1K 7P4 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): VÉZINA, Louis-Philippe [CA/CA]; 206, Route 138, Neuville, Québec G0A 2R0 (CA). D'AOUST, Marc-André [CA/CA]; 939, avenue Manrèse, App. 2, Québec, Québec G1S 2W9 (CA).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— Without international search report and to be republished upon receipt of that report.

[Continued on next page]

(54) Title: METHOD FOR REGULATING TRANSCRIPTION OF FOREIGN GENES



(57) Abstract: The present invention relates to a method of regulating the transcription of transgene in genetically-modified organisms. More specifically, the invention relates to the use of expression vectors harboring the coding sequence of a gene of interest under the transcriptional control of promoting sequences for which activity is regulated by the presence of nitrogen.



*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## **METHOD FOR REGULATING TRANSCRIPTION OF FOREIGN GENES**

### **BACKGROUND OF THE INVENTION**

#### **(a) Field of the Invention**

5           The invention relates to a method of regulating the transcription of transgene in genetically-modified organisms. More specifically, the invention relates to the use of expression vectors harboring the coding sequence of a gene of interest under the transcriptional control of promoting sequences for which activity is regulated by the presence of  
10   nitrogen. Preferably, these constructs are used in transgenic leguminous plants (for example soybean, alfalfa, clover, birdsfoot trefoil, beans, peas, peanuts) where growth is not impaired by lack of mineral nitrogen, and in which induction of expression could be performed at any given time during development, through the addition of a suitable nitrogen source. In a  
15   broader perspective, the invention could be used to induce expression of any given transgene through the addition of any nitrogen source, provided that the organism can be grown adequately in the absence of this nitrogen inducer; as an example within the plant kingdom, duckweed (*Lemna minor*) can adapt to grow either on nitrate or ammonium as nitrogen  
20   source; transgenic duckweed could therefore be grown on nitrate as a sole nitrogen source and expression of the transgene triggered by the addition of ammonium, provided that the cassette contains a promoter from a native gene for which expression is turned on by the addition of ammonium. The invention therefore provides a means of regulating the  
25   expression of a transgenic trait in any organism through the addition of various nitrogenous inducer.

#### **(b) Description of Prior Art**

          Nitrogen is a molecule essential to life. All living organism need nitrogen in order to synthesize amino acids, the building blocks of  
30   proteins, and nucleotides, the building blocks of nucleic acids. It is Ammonium nitrate is the preferred form of mineral nitrogen provided to crops in the form of fertilizer. Nitrate-nitrogen is first reduced to nitrite and then to ammonium through the activity of a metabolic pathway common to most herbaceous plants. Depending on the species, part or all of the  
35   absorbed nitrate will move to leaf cells through the xylem before it is

reduced to ammonium. Ammonium, or other reduced forms of nitrogen are also absorbed (although usually at lower rates) by the root system but their assimilation does not require reduction. These newly absorbed ammonium or ammonium-containing molecules join the endogenous  
5 pools in the cells which is formed by ammonium cycling through amino acids and other nitrogenous molecules. Some species do not metabolize nitrate-nitrogen easily and therefore cannot rely on nitrate as sole nitrogen source; many coniferous species fall into this latter category. Legumes and other symbiotic plant species form a third large class of nitrogen user  
10 within the plant kingdom; they form a metabolic alliance with a microbial organism through which they can fix gaseous nitrogen. This reduced nitrogen is used efficiently by the plant for growth, and therefore, these crops can develop independently of the availability of mineral nitrogen in the soil.

15 Many microbes and wild plant species will adapt extensively to availability of nitrogen sources and can therefore complete their life cycle in the absence of one molecular form of nitrogen, which they could use exclusively and efficiently if available in another growing environment. As for most assimilatory pathways, nitrogen assimilation is tightly regulated in  
20 cells. As an example, the expression of genes encoding nitrate reductase (NaR) and nitrite reductase (NiR), which are responsible for the reduction of nitrate to ammonium, has been extensively described in various microbial and plant species (for a review, see Miflin and Lea, Books 5 and 12, in *The Biochemistry of plants*). Although nitrate is not the only  
25 regulatory molecule involved in the control of NaR and NiR expression, its presence is essential to initiate the cascade of transduction that eventually leads to sustained transcription and translation of these genes. It has been shown that expression of NaR and NiR genes is repressed in leguminous plants when they are grown in the absence of mineral  
30 nitrogen

NiR promoters have been characterized in some plant species (Back et al., 1991, *Plant Molecular Biology* 17:9-18; Sander et al., 1995, *Plant Molecular Biology* 27:165-177). Inducibility of these promoters have also been characterized using marker genes in transgenic plants, where it

was shown that availability of nitrate is required for full activation of transcription.

Assimilatory pathways for other nitrogen sources have also been described, and promoters for genes involved in some of these pathways have also been characterized.

Genetic transformation of microbes have been used for more than 15 years to produce useful recombinant molecules, and applications in the pharmaceutical, cosmaceutical and dermaceutical industries are being currently exploited. This technology has expanded from microbes to plants and animals in the last ten years with the development of techniques required to adapt this general concept to complex eukaryotic organisms. Basically a gene encoding for a protein of interest or a gene encoding for an enzyme responsible for a modification of a metabolic pathway that leads to a molecule of interest, is linked in an appropriate fashion to cis-and trans-acting regulatory sequences, and transferred to a target cell where it is incorporated in the molecular machinery (in a transitory or stable fashion). The transgenic cell, or a tissue or organism regenerated from the transgenic cell will then perform transcription and translation of the transgene and therefore be enabled to accumulate the protein of interest or to perform the new metabolic reaction through the activity of the enzyme of interest.

The emerging industry of molecular farming (production of recombinant molecules in animals or crops) is one of the most promising industry of the coming century. Its promise is to provide safe and renewable molecule factories for the industry. Among the applications that are currently developed are the production of low-cost monoclonal antibodies for therapeutic and diagnostic uses, the production of unlimited amounts of hormones, cytokines and other bio-active molecules for the treatment of chronicle or lethal diseases, the production of bio-safe substitutes for various blood components, the production of unlimited amounts of processing enzymes for the food and pulp industry, the production of low-cost enzymes for waste treatments, and the production of safe bio-active molecules for the cosmetic industry.

Limitations to the application of this technology has often come from the inability of transgenic organisms to accumulate adequate

amounts of the recombinant product, as a result of low transcription rates, improper splicing of the messenger, instability of the foreign mRNA, poor translation rates, hyper-susceptibility of the recombinant protein to the action of endogenous proteases or hyper-susceptibility of the recombinant organism to the foreign protein which result in improper and limited growth or in the worst cases, in strong deleterious effects to the host organism. Inadequacy of production level has a direct impact on the development of applications when profit margins are narrow, or when treatment and/or disposal of residual matter causes bio-safety or environmental problems.

Improvement of the accumulation level of the desired recombinant product thus appears to be one critical factor that warrants commercialization of many applications of molecular farming.

The use of inducible promoters has been proposed, and in some instances used successfully, to counteract the combined effect of all the above-mentioned factors. Strong inducible promoters may succeed in generating high ephemeral transcription rates which result in high transitory accumulation of foreign mRNA and translational product. As a result, when inducibility of expression is paired with adequate synchronized protein recovery procedures, the yield per unit obtained is higher than with the use of constitutive expression.

Several expression cassettes harboring inducible promoters have been developed for microbial production systems, and some are currently available for research purposes. Some inducible promoters are currently used in plant (wound inducibility) or animal (specificity to cells of the mammary glands, PPL) systems, although none reported are using low-cost and bio-safe chemical inducers such as nitrate salts.

It would be highly desirable to be provided with a method of regulating the transcription of transgene in genetically-modified organisms.

#### **SUMMARY OF THE INVENTION**

One aim of the present invention is to provide a method of regulating the transcription of transgene in genetically-modified organisms.

Another aim of the present invention is to provide the use of expression vectors harboring the coding sequence of a gene of interest under the transcriptional control of promoting sequences for which activity is regulated by the presence of nitrogen.

5           The present invention relates to the use of a nitrogen-inducible expression cassettes for the controlled expression of foreign genes in plants. It will be shown from the following description that isolating such a regulatory sequences can be performed so that when cis-acting sequences are appropriately associated to the open reading frame of a  
10 gene of interest, its transcription can be controlled by the addition of specific nitrogen sources.

          In one aspect of this invention, the targeted system uses leguminous plant species, so that constructs containing a nitrate-inducible promoter will be maintained transcriptionally low throughout the growth  
15 period if the transgenic plant is maintained on a nitrate-free medium, thus allowing the development of the plant biomass without interference from the transgenic trait. Upon addition of nitrate to the growth medium, transcription will be induced in a relatively large proportion of the biomass over the following days. Optimization of induction time and protein  
20 accumulation will then be performed in order to maximize recovery of the desired recombinant product.

          Although the following description will make clear that this invention can be easily adapted to nitrate induction on nitrate-deprived transgenic leguminous plants, it should be remembered that this general  
25 concept can also be applied to the development of other production systems, making profit of the wide variety of nitrogen assimilation systems in the microbial, plant and animal kingdoms.

          In one other aspect of this invention, nitrogen inducibility can also be used to maximize protein production in organisms which do not  
30 perform nitrogen fixation through symbiotic association, but that can use variable sources of nitrogen (reduced or oxidized) for growth, and thus possess the ability to develop adequately while one of their nitrogen assimilation pathway is inactive due to lack of one nitrogenous substrate in the growing environment. Using an expression cassettes that controls  
35 the transcription of any gene in this inactive pathway in order to drive the

expression of a gene of interest in such an organism, will allow for inducible expression of the transgenic trait upon addition of the previously lacking nitrogenous compound. As an example, duckweed is a plant species that can grow alternately on nitrate or ammonium; this invention  
5 could be used to develop an expression cassette harboring an ammonium-inducible promoter appropriately linked to a gene of interest so that the induction would be performed on nitrate-grown transgenic duckweed plants.

10 In accordance with the present invention there is provided a method for regulating transcription of a foreign gene in transgenic organisms, comprising the steps of:

- 15 a) preparing a transgenic organism using an expression construct consisting of at least a nitrogen-inducible promoter having a sequence selected from the group consisting of SEQ ID NOS: 1 to 13 and functional fragments and derivatives thereof, and an ORF of a gene, wherein said promoter is operationally located with respect to said gene for expression of said gene.

20 In accordance with a preferred embodiment of the present invention, the method of may further comprise the step of regulating transcriptional expression of said gene by addition or removal of a nitrogen inducer.

In accordance with a preferred embodiment of the present invention, there is provided a method for regulating transcription of a foreign gene in transgenic organisms comprising:

- 25 a) preparing an expression construct consisting of at least a nitrogen-inducible promoter with or without cis-acting sequence, an ORF of a gene, and a polyadenylation signal end site at the 3'end of said construct, wherein said promoter is operationally located with respect to said gene for expression of said gene and modulated for transcriptional expression of said gene by  
30 addition or removal of a nitrogen inducer;
- b) sub-cloning the construct of step a) into a suitable transfection vector for said organism;
- c) transferring said vector into DNA of said organism or a cell  
35 thereof; and



- d) selecting for transgenicity on a suitable medium.

In accordance with a preferred embodiment of the present invention, the method of may further comprise the steps of:

- 5 e) introducing the vector into a suitable *Agrobacterium tumefaciens* strain;  
f) using the *Agrobacterium* strain of step a) to transfer T-DNA into a plant cell;  
g) selecting for transgenicity of said plant cell on a suitable medium;  
10 h) regenerating embryos or plantlets from said transgenic cells; and  
i) growing mature plants from said regenerated embryos.

15 In accordance with a preferred embodiment of the present invention, the cis-acting sequence may be isolated from 5' upstream region of an expressed Nir gene in alfalfa.

In accordance with a preferred embodiment of the method of the present invention, the promoter has the sequence set forth in SEQ ID NO:1 to 13 and functional fragments and derivatives thereof.

20 In accordance with a preferred embodiment of the present invention, the organism is a plant, more preferably a dicotyledonous plant.

In accordance with a preferred embodiment of the present invention, the organism is alfalfa or tobacco.

25 In accordance with a preferred embodiment of the present invention, the nitrogen inducer is nitrate.

In accordance with a preferred embodiment of the present invention, the DNA transfer method is any suitable transfer method including DNA bombardment, electroporation, PEG-mediated DNA transfer and whiskers, among others.

30 In accordance with a preferred embodiment of the present invention, the expression construct comprises at least a nitrogen-inducible promoter and at least one cis- or trans-acting elements.

35 In accordance with a preferred embodiment of the present invention, the organism is a plant, a fungus, a bacteria, a yeast or an animal.

In accordance with a preferred embodiment of the present invention, the promoter or cis-acting sequence is isolated from the 5' upstream region of any gene involved in a nitrogen assimilatory pathway.

5 In accordance with a preferred embodiment of the present invention, the promoter or cis-acting sequence is isolated from the 5'upstream region of any gene for which transcription is modulated by availability of a given nitrogen source.

10 In accordance with a preferred embodiment of the present invention, the promoting or cis-acting sequence is any sequence for which transcriptional activity is regulated by addition or removal of any nitrogen source in or from any living organism's environment.

In accordance with a preferred embodiment of the present invention, the organism from which the promoter or cis-acting sequence is isolated from is any plant, fungus, yeast, bacteria or animal.

15 In accordance with a preferred embodiment of the present invention, there is provided a promoter for promoting transcription of a foreign gene in transgenic organisms, which comprises a nitrogen-inducible promoter with or without cis-acting sequence for expression of said gene and adapted to be modulated for transcriptional expression of  
20 said gene by addition or removal of a nitrogen inducer.

Preferably, the promoter has a sequence selected from the group consisting of SEQ ID NOS: 1 to 13 and functional fragments and derivatives thereof.

25 In accordance with a preferred embodiment of the present invention, there is provided a terminator allowing expression of a foreign gene in transgenic organisms being used in combination with a promoter, which comprises a polyadenylation signal end site for insertion at the 3'end of said gene, wherein said terminator is operationally located with respect to said gene and said promoter and thereby allows expression of  
30 said gene.

Preferably, the terminator has a sequence selected from the group consisting of SEQ ID NOS: 14 to 16 and functional fragments and derivatives thereof.

35 For the purpose of the present invention the following terms are defined below.

The expression "functional fragments or derivatives thereof" is intended to mean any derivative or fragment of sequences SEQ. ID. NOS:1-16 which allow for an equivalent level of expression of a foreign gene as the promoter of the present invention set forth in SEQ. ID. NOS:1-13 or as the terminator of the present invention set forth in SEQ. ID. NOS:14-16.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 illustrates GUS expression level using promoter Nir (SEQ ID NOS:2-13) and terminator NOS in the leaves of transgenic tobacco plants before (empty columns) and after (filled columns) nitrate fertilization of the plants. Tobacco plants were transformed with full-length and deletions of the alfalfa NiR promoter and NOS terminator functionally positioned to control transcription and termination of the GUS reporter gene, as described in Methods. GUS activity was measured according to Jefferson et al. (1987, *EMBO J.* 13:3901-3907) prior and after induction by nitrate.

Fig. 2 illustrates GUS expression level using promoter Nir (SEQ ID NOS:2, 3, 5-6) and terminator Nir (SEQ ID NOS:15-16) in the leaves of transgenic tobacco plants before (empty columns) and after (filled columns) nitrate fertilization of the plants. Tobacco plants were transformed with full-length and deletions of the alfalfa NiR promoter and NiR terminator functionally positioned to control transcription and termination of the GUS reporter gene, as described in Materials and Methods below. GUS activity was measured according to Jefferson et al (1987, *EMBO J.* 13:3901-3907) prior and after induction by nitrate.

Fig. 3 illustrates GUS expression level using promoter NiR and terminator NOS in the leaves of nodulated transgenic alfalfa plants before (empty columns) and after (filled columns) nitrate fertilization. Alfalfa genotype 11.9 was transformed with constructs GC2-E, GC2-B, DC1-D and 35S functionally positioned to drive expression of reporter gene GUS, as described in Khoudi et al (1997, *Gene* 197:343-351). Following regeneration, transgenic plants were transferred to sterile vermiculite and inoculated with *Rhizobium* strain Balzac (Nitragin). Plants were allowed to grow for 3 weeks with repeated additions of nitrate-free Hoagland's

solution; GUS activity was then measured in first fully expanded leaves as described by Jefferson et al. (1987, *EMBO J.* 13:3901-3907). Plants were then fertilized with 40 mM nitrate for two days. GUS activity was again measured in first fully-expanded leaves. Data shown herein are ratios  
5 between post- and pre-nitrate induction GUS specific activities.

### **DETAILED DESCRIPTION OF THE INVENTION**

Following is a detailed description of the method used to generate transgenic tobacco and alfalfa lines that can be modulated in  
10 their expression of a reporter gene. It should be remembered that variations could be brought to the method by which nitrogen-inducible promoters could be isolated, by which they could be linked to ORFs in the construct used for expression in plants, by which different cis- and trans-acting elements of the constructs are used and spatially arranged, by  
15 which the inducibility by nitrogen is demonstrated and used, while remaining within the scope of this invention.

In this embodiment, a NiR cDNA strand was first isolated from alfalfa using RT-PCR with primers deduced from a consensus plant NiR sequence. This cDNA stretch was then used either to perform  
20 upstream/downstream genome walking. The NiR promoter region and deletions, the 5'UTR and the NiR terminator were then positioned functionally to control transcription and terminaison of reporter gene GUS. These constructs were inserted into suitable expression vectors for DNA bombardment onto tobacco and alfalfa leaves and for *Agrobacterium*  
25 mediated DNA transfer as described by Desgagnés et al. (1995, *Plant Cell Tissue Organ Cult.* 42:129-140). These two DNA transfer methods were used to demonstrate that expression of the reporter gene can be modulated by addition or removal of nitrate in the growing medium.

### **Materials and Methods**

#### **Biological material**

*E. coli* strain DH5- $\alpha$  was used to perform all cloning steps. Cold resistant alfalfa genotype 11.9 was used for all experiments including

stable transformation using *A. tumefaciens* infection (Desgagnés et al. (1995, *Plant Cell Tissue Organ Cult.* 42:129-140).

#### Isolation of total RNA

5           Total RNA was extracted using a hot phenol method essentially as described by de Vries et al. (1988, B6 page 1, In: Gevin SB and Shilperoot RA editors, *Plant Molecular Biology Manual*, Dordrecht: Kluwer Academic Publisher).

#### 10   RT-PCR

RT-PCR was used to produce a DNA fragment corresponding to one abundant NiR mRNA molecular species from leaf total mRNA. A conserved region was first identified from 5 public plant NiR ORFs, namely Genbank sequences #AB006032 (*Arabidopsis* Nir mRNA), #  
15   X66145 (*Tobacco* partial Nir mRNA), #U10419 (*Bean* complete Nir cds), #X07568 (*Spinach* Nir mRNA), and #U90429 (*Glycine max* Nir complete cds). Degenerated oligonucleotides were deduced from two conserved regions, namely Nir5 - 5' GATATTGATGTTAGACTCAAGTGGC 3' (SEQ ID NO:17), at the 5' end and Nir3 - 5' CACYSATTCCACTTCCTWGWC 3'  
20   (SEQ ID NO:18), at the 3' end of the coding strand. A reverse transcription reaction was first performed with 200 units of M-MLV reverse transcriptase (RT) for 1 hour at 37°C using 1 µg of total leaf RNA, 4 mM dNTP (1 mM each), 5 µM random hexamer primers in a 1X M-MLV-RT buffer (50 mM Tris-HCl pH 8.3, 75 mM KCl, 3mM MgCl<sub>2</sub>). The PCR  
25   reaction was performed in a Perkin Elmer Cetus GenAmp PCR system 9600 (EG&G, Wellesley, MA), using 2.5 units of Taq DNA polymerase, 2 µM Nir5 primer, 2 µM Nir3 primer, 800 µM dNTPs (200 µM each) in a 1X PCR buffer (20 mM Tris-HCl pH 8.4, 50 mM KCl, 2 mM MgCl<sub>2</sub>). The cycling program used was: an initial 4 min at 94°C, 30 cycles of 1 min at  
30   94°C, 30 sec at 55°C, and 3 min at 72°C. An extension period of 7 min at 72°C was included in the program.

**DNA sequencing**

DNA sequencing was performed as described by Sanger et al. (1977, *P.N.A.S. USA*, 74:5643-5647).

5

**Genome walking**

Walking upstream of the Alfalfa Nir cDNA fragment cloned from the RT-PCR reaction was performed using the Universal Genome Walker Kit from Clontech Laboratories (Palo Alto, CA) (Cat. #1807-1). The NiR-specific custom primers used to amplify sequences upstream of the coding sequence were:

10

Nir1106r - 5' TTGTCACATCAGCACATCCGTCTTTGC 3' (SEQ ID NO:19), and

15

Nir1061r - 5' TCGCCAAGTATCTTGTTGAGCACTTG 3' (SEQ ID NO:20).

The amplified 3775 bp fragment was subcloned into pGEM-T Easy vector (Promega, Madison, WI) (Cat #A1360) for further analysis. The resulting plasmid was named pGNir4c.

The downstream walking was performed as the upstream walking using the following NiR specific primers:

20

Nir1c - 5' ATGTCTTCCTTCTCAGTACGTTTCCTC 3' (SEQ ID NO:28), and

Nir138c - 5' CAAGTTGATGCATCAAGGTGGGAGCCTAGA 3' (SEQ ID NO:29).

25

The amplified 3508 bp fragment was subcloned into pGEM-T easy vector (Promega, Madison, WI) (Cat #A1360) for further analysis. The resulting plasmid was named pGN3'1.

**Construction of expression cassettes and vectors**

The cassettes for expression analysis using the GUS reporter gene were assembled as follows. A promoterless GUS cassette was digested from pBI101 with HindIII and EcoRI, and was inserted into the HindIII and EcoRI sites of the pUC19 polycloning site. The resulting plasmid was named pBI201 and was used for further constructs. The Nir upstream sequences were PCR amplified using the AP2 primer from the Universal Genome Walking Kit as upstream positioned primer, and either one of custom-designed downstream primers ending with a SmaI

35

restriction site. The 4 primers were positioned either in the 5' UTR region of the gene (Nir-23r-Sma-5'AGAGCCCGGGAGAAGAGAGTGTGTTTG3' (SEQ ID NO:21)), at the end of the transit peptide coding sequence (Nir51r-Sma - 5' TTCTCCCGGGGGACGAGAGATGGATGGT 3' (SEQ ID NO:22)), 50 bp after the transit peptide coding sequence (Nir103r-Sma - 5' TTCTCCCGGGGTTGAA-ACAGGTGCAACTGA 3' (SEQ ID NO:23)), and 100 pb after the transit peptide coding sequence (Nir158r-Sma - 5' TTCTCCCGGGTAACCATCTTTTCCTCA 3' (SEQ ID NO:24))  
Amplification was performed under standard conditions with pGNir4c plasmid as template.

The amplified fragments were digested with specific restriction enzymes in order to produce 5' deletions of the Nir promoter. The pNir3k-23 was produced by digesting the fragment previously amplified by AP2 and Nir-23r-Sma primers with XmaI, and inserting the resulting fragment into pBI201 previously digested with XmaI. A similar strategy was used to produce the pNir3k51, pNir3k103, and pNir3k158 plasmids except that the downstream primers used were Nir51r-Sma, Nir103r-Sma, and Nir158r-Sma, respectively. The pNir2.2k-23 was produced from a SmaI-BglII digestion of the AP2 - Nir-23-Sma amplified fragment inserted into the pBI201 previously digested with SmaI and BamHI. The same strategy was used to produce the pNir2.2k51, pNir2.2k103, and pNir2.2k158 plasmids except that the downstream primers used were Nir51r-Sma, Nir103r-Sma, and Nir158r-Sma, respectively. Fidelity and orientation of the insertions were verified by digestion with restriction enzymes. These deletion fragments were ligated to the 5'terminus of the GUS reporter gene in pBI201, and used for transitory expression studies using DNA bombardment. Upon identification of the adequate deletion fragments, they were sub-cloned into a binary plant expression vector such as pBI101 (Clonetech).

For the construction of the cassettes containing the NiR terminator downstream of the GUS gene in addition to the NiR promoter, the following NiR specific primers were used:

Nir2514c-Sac - 5' AGAAGAGCTCAGTATATAGGTATTTGGTGA 3'(SEQ ID NO:30)

Nir2728c-Sac - 5' AGAAGAGCTCTTGTACATTTGGATAAGTCA 3' (SEQ ID NO:31)

Nir3029r-Eco - 5' AGAAGAATTCGTTTTCCCGATACTTCAACT 3' (SEQ ID NO:32)

5 A 617 bp terminator fragment was PCR amplified using the primers Nir2514c-Sac and Nir3029r-Eco, and a 503 bp terminator fragment was PCR amplified using the primers Nir2728c-Sac and Nir3029r-Eco. The fragments obtained were digested with SacI and EcoRI and inserted into the plasmids containing the NiR-GUS constructs after  
10 deletion of the NOS terminator between the SacI and EcoRI sites.

These recombinant plasmids were used for stable integration through *A. tumefaciens* infection as described below.

#### 15 ***Agrobacterium*-mediated DNA transfer and regeneration of transgenic alfalfa lines**

The recombinant plasmids were introduced into *Agrobacterium tumefaciens* strain LBA4404 by electroporation as described in Khoudi et al (1997, *Gene* 197:343-351). Selected *Agrobacterium* strains were then co-cultivated with leaf disks from genotype C5-1 for 4 days in the absence  
20 of selection pressure (kanamycin). Following this incubation period, leaf disks were washed and pampered, and then allowed to form calli onto medium B5H. Calli were then transferred for 21 days on SH medium for embryo induction and for 28 days on BOi2Y for embryo development. Torpedo-shaped embryos were removed from BOi2Y and placed on MS  
25 medium for regeneration. Kanamycin was present in all cultivation medium except for co-cultivation and regeneration on MS. This method is described in length in Desgagnés et al (1995, *Plant Cell Tissue Organ Cult.* 42:129-140). Rooted plantlets were grown to maturity in the greenhouse. Integration of the transgene was verified by PCR  
30 amplification of a NiR-GUS fragment from genomic DNA. The primers used were:

Nir-102c – 5' CACACTTCTTCACTCACCTCTCAA 3' (SEQ ID NO:25)

Nir-2016c – 5' ATCTAGGAGGGGCAGACATTG 3' (SEQ ID NO:26)

GUS228r – 5' TCGGTATAAAGACTTCGCGCTGAT 3' (SEQ ID NO:27)

35



### ***Agrobacterium*-mediated DNA transfer and regeneration of transgenic tobacco lines**

The recombinant plasmids were introduced into *Agrobacterium tumefaciens* strain LBA4404 by electroporation as described in Khoudi et al (1997, *Gene* 197:343-351). Selected strains were co-cultivated with leaf disks for 2 days on MS medium without kanamycin. After this period, the explants were transferred to the selection medium (MS with Kanamycin). The explants were kept on this medium for 3 weeks to allow the formation of calli and shoots from the transfected cells. The kanamycin resistant shoots were transferred into the rooting MS medium. Rooted plantlets were grown to maturity in the greenhouse. Integration of the transgene was verified by PCR amplification of a NiR-GUS fragment from genomic DNA. The primer used were:

- Nir-102c – 5' CACACTTCTTCACTCACCTCTCAA 3' (SEQ ID NO:25)  
Nir-2016c – 5' ATCTAGGAGGGGCAGACATTG 3' (SEQ ID NO:26)  
GUS228r – 5' TCGGTATAAAGACTTCGCGCTGAT 3' (SEQ ID NO:27)

### **Nitrate induction**

Transgenic and non-transgenic tobacco and alfalfa plants were grown in vermiculite medium without nitrate. Mineral balance was kept by repeated additions of nitrate-free Hoagland's solutions (Hoagland and Varnon, 1950, Circular 347, *California Agr. Exp. Stat. Berkeley*). Nitrate induction was performed by watering the plants with 20-20-20 fertilizer at a concentration of 5gL<sup>-1</sup> or as an alternative with Hoagland's solution supplemented with 40 mM nitrate.

### **NiR promoter activity in tobacco leaves**

The NiR derived promoters were placed upstream of the GUS reporter gene in transcriptional and translational fusions. The 5' deletions of the NiR promoter analyzed here consisted in (1) a putative full length promoter comprising 2813 bp upstream of the initial ATG of the coding region, (2) a 1905 bp version of the promoter, and (3) a shorter 1111 bp version of the promoter. The 3' end of the promoter was fused to the 5' end of the GUS coding region to form transcriptional and translational

fusions. Translational fusions analyzed allowed the production of  $\beta$ -glucuronidase containing (1) the NiR chloroplast transit peptide, (2) the NiR transit peptide with an additional 17 amino acids from NiR, and (3) the NiR transit peptide with an additional 36 amino acids from NiR. The twelve combinations of 5' and 3' deletions of the NiR promoter introduced into tobacco plants are presented in Fig. 1.

The gene constructs were transferred into tobacco plants using the *Agrobacterium*-mediated transfection method (Khouidi et al., 1997, *Gene* 197:343-351). Transgenic plants were transferred to growth chambers and analyzed for their leaf  $\beta$ -glucuronidase content before and after nitrate fertilization. Fig. 1 presents the median level of  $\beta$ -glucuronidase activity measured in the 1<sup>st</sup> expanded leaf of plantlets.

All the NiR derived promoters showed reactivity to nitrate induction. Between 5 and 10 fold increase of  $\beta$ -glucuronidase expression was generally observed, irrespectively of the promoter truncation, indicating that important nitrate responsive elements are contained within the first 1.1 kb upstream of the initial ATG. Both 5' and 3' deletions of the NiR promoter led to important modifications of  $\beta$ -glucuronidase activity. The highest level of GUS expression was obtained with the 2.8 kb promoter, indicating that the far upstream regions have a regulatory role for the level of NiR expression in the leaves.

The translational fusions of the promoter to the GUS coding region resulted in variable expression level depending on the extension of the 5' end of the promoter. However, the shortest fusion (containing the 17 a.a. NiR transit peptide fused to the amino-terminal end of the  $\beta$ -glucuronidase) constantly resulted in the highest level of activity for all three 5' end truncations. This short translational fusion, combined with the longest extension of upstream promoter regions gave rise to the strongest promoter (3kb+50). When induced, this specific construct resulted in more than 13-fold the level of GUS expression obtained with the 35S-GUS-NOS construct.

When considering the longest 5' extension of the promoter, the transcriptional fusion to the GUS gene (3kb-5) was ~7 times less effective than the short translational fusion (3kb+50). However, in it's induced state, the level of GUS expression in the plants harboring the 3kb-5 promoter deletion was more than 1,8-fold that observed in the 35S-GUS-NOS plants.

Taken together, the results presented here clearly indicate that the sequences upstream of the alfalfa NiR gene have the capacity to drive high and inducible expression of an exogenous gene in tobacco leaves.

10

#### **Efficiency of NiR terminator**

Tobacco plants were transformed with constructs consisting of promoter NiR and deletions (SEQ ID NOS: 2, 3, 5 and 6), and 35S, together with 3'UTR sequences and terminator (SEQ ID NOS: 15 and 16), functionally positioned to drive transcription and termination of reporter gene GUS. Growth, nitrate induction and GUS activity measurements were performed as per experiment illustrated in Fig 1. Results shown in Fig 2 demonstrate that the terminator sequence of Nir allows termination of transcription into a translatable messenger RNA.

20

#### **NiR promoter activity in alfalfa leaves**

Transgenic alfalfa plant containing the gene constructs presented in Fig. 3 were obtained using the *Agrobacterium*-mediated transfection method of Desgagnés et al. (1995, *Plant Cell Tissue Organ Cult.* 42:129-140). The *in vitro* plants were transferred into growth chamber to allow a normal vegetative growth. Cuttings from each transgenic line were grown in vermiculite and fertilized with nitrate-free Hoagland medium. After two weeks, the roots were inoculated with Nitragin (LiphaTech inc., Milwaukee, WI). Two weeks after inoculation, nodules had developed on the roots. Nodulated plants were allowed to continue their vegetative growth for at least another week before the fluorometric measurement of  $\beta$ -glucuronidase activity (before induction)

30

was performed. After the measurement, the plants were watered with Hoagland medium containing 40 mmol KNO<sub>3</sub>. Two days after induction, leaf  $\beta$ -glucuronidase activity was measured to evaluate the nitrate inducibility of the NiR promoter in alfalfa leaves. Results are presented in

5 Fig. 3. Results show that promoter NiR induces expression of GUS reporter gene upon addition of nitrate in nodulated alfalfa plants. Taken together, this last series of result demonstrate that NiR promoter inducibility can be used to positively regulate expression of a foreign gene in alfalfa plants when fixation of atmospheric nitrogen is replaced by

10 nitrate assimilation.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the

15 principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

20

**WHAT IS CLAIMED IS:**

1. A method for regulating transcription of a foreign gene in transgenic organisms, comprising the steps of:
  - a) preparing a transgenic organism using an expression construct consisting of at least a nitrogen-inducible promoter having a sequence selected from the group consisting of SEQ ID NOS: 1 to 13 and functional fragments and derivatives thereof, and an ORF of a gene, wherein said promoter is operationally located with respect to said gene for expression of said gene.
2. The method of claim 1, which further comprises the step of regulating transcriptional expression of said gene by addition or removal of a nitrogen inducer.
3. A method for regulating transcription of a foreign gene in transgenic organisms comprising:
  - a) preparing an expression construct consisting of at least a nitrogen-inducible promoter with or without cis-acting sequence, an ORF of a gene, and a polyadenylation signal end site at the 3'end of said construct, wherein said promoter is operationally located with respect to said gene for expression of said gene and modulated for transcriptional expression of said gene by addition or removal of a nitrogen inducer;
  - b) sub-cloning the construct of step a) into a suitable transfection vector for said organism;
  - c) transferring said vector into DNA of said organism or a cell thereof; and
  - d) selecting for transgenicity on a suitable medium.
4. The method of claim 3, wherein said organism is a plant.
5. The method of claim 4, which further comprises the steps of:
  - e) introducing the vector into a suitable *Agrobacterium tumefaciens* strain;

- f) using the *Agrobacterium* strain of step a) to transfer T-DNA into a plant cell;
  - g) selecting for transgenicity of said plant cell on a suitable medium;
  - h) regenerating embryos or plantlets from said transgenic cells; and
  - i) growing mature plants from said regenerated embryos.
6. The method of claim 3, wherein the cis-acting sequence is isolated from 5' upstream region of an expressed Nir gene in alfalfa.
7. The method of claim 3, wherein the promoter has the sequence set forth in SEQ ID NOS:1 to 13 and functional fragments and derivatives thereof.
8. The method of claim 1, wherein the organism is selected from the group consisting of a plant, a fungus, a bacteria, a yeast and an animal.
9. The method of claim 8, wherein the plant is a dicot.
10. The method of claim 9, wherein the dicot is alfalfa or tobacco.
11. The method of claim 2, wherein the nitrogen inducer is nitrate.
12. The method of claim 3, wherein the DNA transfer method is any suitable transfer method including DNA bombardment, electroporation, PEG-mediated DNA transfer and whiskers.
13. The method of claim 3, wherein the expression construct comprises at least a nitrogen-inducible promoter and at least one cis- or trans-acting elements.

14. The method of claim 3, wherein the promoter or cis-acting sequence is isolated from the 5' upstream region of any gene involved in a nitrogen assimilatory pathway.

15. The method of claim 3, wherein the promoter or cis-acting sequence is isolated from the 5'upstream region of any gene for which transcription is modulated by availability of a nitrogen source.

16. The method of claim 3, wherein the promoting or cis-acting sequence is any sequence for which transcriptional activity is modulated by addition or removal of any nitrogen source in or from any living organism's environment.

17. The method of claim 3, wherein the organism from which the promoter or cis-acting sequence is isolated from is any plant, fungus, yeast, bacteria or animal.

18. A promoter for promoting transcription of a foreign gene in transgenic organisms, which comprises a nitrogen-inducible promoter with or without cis-acting sequence for expression of said gene and adapted to be modulated for transcriptional expression of said gene by addition or removal of a nitrogen inducer.


19. The promoter of claim 18, wherein said promoter has a sequence selected from the group consisting of SEQ ID NOS: 1 to 13 and functional fragments and derivatives thereof.

20. A terminator allowing expression of a foreign gene in transgenic organisms being used in combination with a promoter, which comprises a polyadenylation signal end site for insertion at the 3'end of said gene, wherein said terminator is operationally located with respect to said gene and said promoter and thereby allows expression of said gene.


21. The terminator of claim 19, wherein said terminator has a sequence selected from the group consisting of SEQ ID NOS: 14 to 16 and functional fragments and derivatives thereof.




1 / 3

*Fig. 1*

2 / 3

*Fig. 2*

3 / 3

*Fig. 3*

## SEQUENCE LISTING

<110> VÉZINA, Louis-Philippe  
D'AOUST, Marc-André  
MEDICAGO INC.

<120> METHOD FOR REGULATING TRANSCRIPTION OF  
FOREIGN GENES

<130> 14149-3"PCT"

<150> US 60/157,133

<151> 1999-10-04

<160> 32

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 3714

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating  
expression of foreign genes

<400> 1

ctgtacattc	atcttgccgc	ctttgcattc	acttggccac	aaagagtaga	gagaaggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgaggggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccatttac	ttgagggtata	180
cttattatct	tgatcaataa	atttgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaagt	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggt	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgattgtgt	ttggtcagac	ttcagagtca	gatgactctg	tttgataaaa	cagcttaatt	420
aagcgcttat	agaatatcat	atgattgtgt	ttggtcagac	ttcagagcat	ctcttggttt	480
ctctggcaat	catatgccta	agaaataaat	agtatcata	gatttgtgtt	ggtcagactt	540
cagagtcaga	tgaccctgtt	tgggtaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttattt	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaataata	gctatcctgg	agagcttggt	gaaataacca	gaaaagaact	tatggacacg	720
tcatgagctg	ttacataaag	atctccctaa	cagtcctcaa	agtgtttatg	ccagtagata	780
aattcaaata	agtcaatcta	aacagaccct	aatccatta	tggtagctat	catttttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taacataatg	ataacacatt	attttgacac	900
aatggggcag	atctagcaat	ttaactctgg	agtccttcaa	gactgctgtt	cttacgaaat	960
tcacgtccct	gaatcatgtt	cctgtatgga	agcctgaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaaggtttac	aaaatatacc	ctgcgggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttgagagagc	aatccttggtg	1140
ccaagtttga	agtaattttt	gtgtagcata	tgttgagcta	cctacaattt	acatgatcac	1200
ctagcattag	ctctttcact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtaa	tgctaccctt	actttggcca	actcatcggg	gattttacatt	1320
cagaaaaatat	acatgacttc	aaccatactt	aaaccctttt	ttgtaagata	actgaatgtt	1380
catatttaaat	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttagg	1440
acaacaagat	tgtgggtctg	tactgttatt	tatttatttt	tttttttagca	gaaacacctt	1500
atcttttggt	tcgtttgaat	gtagaatgaa	aataaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagttagga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atttatgtgt	tttttctttt	1680
ttctaccgga	acttttttaga	acggatccca	actcgttccg	gggccgctac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	acccattcac	1800
catttgattc	agatgtgggtg	actagagata	aagcatacta	atttgactct	tggaaaccca	1860
taaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920

tgtgggtttgg	tttccattct	gatttatgcg	gcgacttgta	attttaaact	taggaggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagtg	tcactttgat	agttcaattt	2160
catcttggtc	tcttattcct	tttataattc	taattcttct	tgtgtaaact	atttcatgta	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcctc	actaactcaa	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacaa	aattttacatc	aaatttcgac	atcgttttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaacta	agttacccta	caacttaaag	aacttaaaga	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	taatgttgac	tttgacaccg	2580
caccacatgt	gcctttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgctctatt	2640
ataaggcgac	ccttagctcc	aaccaaaagg	cgatggagtt	aagaaagaaa	ctcttgctta	2700
cttgtaaggt	ccacacttct	tcactcacct	ctcaattttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttctca	ccccaccatc	catctctcgt	cccaacaaaa	catggctact	2880
atctgctgca	actccatcag	ttgcacctgt	ttcaacacca	caagttgatg	catcaagggt	2940
ggagcctaga	gttgaggaaa	aagatgggta	ctgggttttg	aaggaagagt	atagaggggg	3000
tattaatcct	caggagaaa	ttaagattca	gaaagaacct	atgaagcttt	ttatggaagg	3060
tggtgattaat	gatttggtca	atatgtctct	tgaagagatt	gaaagctcta	agcttactaa	3120
agatgatatt	gatgttagac	ttaaatggct	tggtcttttt	catagaagga	aacatcattg	3180
taagtttttt	taccttcttt	ttatacctca	aagttctctc	atactctgta	tttgttttatt	3240
agttttttgta	gacttaaata	ttctctttga	tttacatagt	gaaactccat	ttttgtttcc	3300
gaaattgtag	tgtgtatagt	ctagaaaatt	aagaagtaga	caaaatgatt	tatgagattg	3360
taaattgtag	gctttttatc	aattttattaa	tttttagagac	caaaatttgc	ctatcttatt	3420
tggaccaata	ttgtatgtca	ggatcgacat	gagtttagta	aaatcatgac	ggcaccatga	3480
ctgtgttgaa	gcttctttgt	gtaactttta	ccaaaattat	atggcacacc	ataattatgc	3540
aaactcaccg	tcgatccaaa	catagaaaatt	cggtgttaat	ctttgtgaga	ataaaaagct	3600
atgagttatg	ttgtactaat	ttattttccat	tgtgaaaatc	agatggtaga	tttatgatga	3660
gactgaaact	tccaaatggg	gtaacaacaa	gtgctcaaac	aagatacttg	gcga	3714

&lt;210&gt; 2

&lt;211&gt; 2808

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 2

ctgtacattc	atcttgccgc	ctttgcattc	acttgggcac	aaagagtaga	gagaaggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgagggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccatttac	ttgaggtata	180
cttattatct	tgatcaataa	atttgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaatg	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggg	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgattgtgt	ttggtcagac	ttcagagtca	gatgactctg	tttggtataa	cagcttaatt	420
aagcgcttat	agaatatcat	atgattgtgt	ttggtcagac	ttcagagcat	ctcttggttt	480
ctctggcaat	catatgccta	agaaataaat	agtatcatat	gattgtgttt	ggcagactt	540
cagagtcaga	tgaccctgtt	tgggtaaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttatct	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaatatata	gctatcctgg	agagcttgtg	gaaataacca	gaaaagaact	tatggacacg	720
tcagagactg	ttacataaag	atctccctaa	cagtctcaaa	agtgtttatg	ccagttagata	780
aattcaataa	agtcaatcta	aacagaccct	aaatccatta	tggtacctat	catttttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taacataatg	ataacacatt	attttgacac	900
aaatgggcag	atctagcaat	ttaactctgg	agtccttcaa	gactgctgtt	cttaacgaagt	960
tcacgtccct	gaatcatgtt	cctgtatgga	agcctgaaag	acctcaaat	ctaaaagggtg	1020
gcgataaatt	gaagggtttac	aaaatatacc	ctgcgggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttgagagac	aatccttgtg	1140

ccaagtttga	agtaattttt	gtgtagcata	tggtgagcta	cctacaattt	acatgatcac	1200
ctagcattag	ctctttcact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtaa	tgcctaccct	actttggcca	actcatcggg	gatttacatt	1320
cagaaaatat	acatgacttc	aaccatactt	aaaccctttt	ttgtaagata	actgaatggt	1380
catatttaat	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttgg	1440
acaacaagat	tgtgggtctg	tactgttatt	tatttatttt	tttttttagca	gaaacacctt	1500
atcttttgtt	tcgtttgaat	gtagaatgaa	aataaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagtttaga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atttattgtg	tttttctttt	1680
ttctaccgga	actttttaga	acggatccca	actcgtttccg	gggcccgtac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	accattcac	1800
catttgattc	agatgtgggtg	actagagata	aagcatacta	atttgactct	tggaaaccca	1860
taaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tggtctctatg	1920
tgtggtttgg	tttccattct	gatttatgcg	gcgacttgta	atttaaaatc	taggaggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatggttc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagttg	tcactttgat	agttcaattt	2160
catcttggtt	tcttattcct	tttataatc	taattcttct	tggtgaaact	atttcatgta	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcctc	actaactcaa	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacia	aatttacatc	aaatttcgac	atcgttttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtta	ttattcatat	atttgaacta	agttacccta	caacttaaa	aacttaaa	2520
actcgggttg	agacctgggg	acgaaaatgt	aatgagactt	taatggtgac	tttgacaccg	2580
caccacatgt	gccttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgctctatt	2640
ataaggcgac	ccttagctcc	aaccaaagga	cgatggagtt	aagaaagaaa	ctcttgctta	2700
cttgtaaggt	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctcc		2808

&lt;210&gt; 3

&lt;211&gt; 2069

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 3

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaa	aagtcaatct	60
aaacagaccc	taaatccatt	atggtaccta	tcatttttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tattttgaca	caaattgggca	gatctagcaa	180
tttaactctg	gagtccttca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240
tcctgtatgg	aagcctgaaa	gacctcaaat	tctaaaaggt	ggcgataaat	tgaagggtta	300
caaaatatac	cctgcgggct	tgacacagag	gcaagctctt	tataaccttc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttggagag	caatccttgt	gccaagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttaactgaga	gaatgaagtt	ttaggaatga	gtatgaccat	ggagtcggca	tggctttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggatttacat	tcagaaaata	tacatgactt	600
caaccatact	taaaccctt	tttgtaagat	aactgaatgt	tcataatttaa	tgttgggttg	660
tagtgttttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttattttatt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagtttag	atcctctccg	gtcaccggaa	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actattattt	tattttattgt	gtttttcttt	tttctaccgg	aacttttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgtaagtttt	cctttacaga	taccattcca	ccatttgatt	cagatgtggg	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaccc	ataaagttta	tggtatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtgggttg	gtttccattc	1200
tgattttatgc	ggcgacttgt	aatttaaaat	ctaggagggg	cagacattga	acaatcccaa	1260

tattttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	gtgactgaga	1320
ttgagtcata	catttcacta	agaaatgggt	ccaagtacca	aactatcatg	accagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcactctggc	ttcttattcc	1440
ttttataatt	ctaattcttc	ttgtgtaaac	tatttcattg	attatttttc	tttaaaattt	1500
acatgtcatt	tattttgcct	cactaactca	atattgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaatttacat	caaatttcga	catcgtttat	tatgttcatt	ggatgattaa	1620
caaatataac	aaactttgca	actaattaac	caccaactga	atataattaa	ctataactgt	1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaaatgta	attattcata	1740
tatttgaact	aagtaccctt	acaacttaaa	gaacttaaa	aactcgggtt	gagacctggg	1800
gacgaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgctttttac	1860
atatagttta	tatgacaagt	aatgacaatc	cttgctctat	tataaggcga	cccttagctc	1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgtaagg	tccacacttc	1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac	2040
aaactcattc	caaacacact	ctcttctcc				2069

&lt;210&gt; 4

&lt;211&gt; 1220

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 4

gatcctctcc	ggtcaccgga	aagtttcagt	agaagaaaca	aaacaccgtg	actaaaatga	60
tactattatt	ttattttattg	tgtttttctt	ttttctaccg	gaacttttta	gaacggatcc	120
caactcgttc	cggggccgct	acaactgaaa	caaaagaaga	tattttctct	ctcttcagaa	180
atgtaagttt	ctctttacag	ataccatttc	accatttgat	tcagatgtgg	tgactagaga	240
taaagcatal	taatttgact	cttggaacc	cataaagttt	atgttatccg	tgttctggac	300
caatccactt	gggggcataa	cctgtgtcta	tggtgtgttt	ggtttccatt	ctgatttatg	360
cggcgacttg	taatttaaaa	tctaggaggg	gcagacattg	aacaatccca	atattttaat	420
aacttatgca	agattttttt	tattaatgag	atgatgtgtt	tgtagctgag	attgagtcac	480
acatttcact	aagaaatggg	tccaagtacc	aaactatcat	gacccagttg	caaacatgac	540
gttcgggagt	ggtcactttg	atagttcaat	ttcatcttgg	cttcttattc	cttttataat	600
tctaattctt	cttggtgaaa	ctatttcatt	tattttttt	ctttaaaatt	tacatgtcat	660
ttattttgcc	tcactaactc	aattttgcat	ataacaatga	taagtgatat	tttgactcac	720
aaaattttaca	tcaaatttcg	acatcgttta	ttatgttcat	tggatgatta	acaaatataa	780
caaactttgc	aactaattaa	ccaccaactg	aatataatta	actataactg	tgaaagtagt	840
taaccatatt	tttttagatgt	atatatcatc	cgttgaaatg	aattattcat	atatttgaac	900
taagttaccc	tacaacttaa	agaacttaaa	gaactcgggt	tgagacctgg	ggacgaaaat	960
gtaatgagac	tttaatgttg	actttgacac	cgcaccacat	gtgcctttta	catatagttt	1020
atatgacaag	taatgacaat	ccttgctcta	ttataaggcg	acccttagct	ccaaccaaag	1080
gacgatggag	ttaagaaaga	aactcttgct	tacttgtaag	gtccacactt	cttcactcac	1140
ctctcaattt	catctacaa	aaatgtccaa	acttctcttt	ctcacaatca	caaactcatt	1200
ccaaacacac	tctcttctcc					1220

&lt;210&gt; 5

&lt;211&gt; 2863

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 5

ctgtacattc	atcttgccgc	ctttgcatte	acttggccac	aaagagtaga	gagaagggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgaggggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccatttac	ttgaggata	180

cttattatttt	tgatcaataa	atttgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaagt	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggg	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgatttgtg	ttgggtcagac	ttcagagtca	gatgactctg	tttggataaa	cagcttaatt	420
aagcgcttat	agaatatcat	atgatttgtg	ttgggtcagac	ttcagagcat	ctcttggttt	480
ctctggcaat	catatgccta	agaaataaat	agtatcatat	gatttgtgtt	gggtcagactt	540
cagagtcaga	tgaccctggt	tgggttaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttattt	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaataata	gctatcctgg	agagcttgtg	gaaataacca	gaaaagaact	tatggacacg	720
tcatgagctg	tttacataag	atctccctaa	cagctcctaa	agtgtttatg	ccagtagata	780
aattcaaata	agtcaatcta	aacagaccct	aaatccatta	tggtagctat	cattttagct	840
tattccatct	ttattaagaa	tgatcatgaga	taacataatg	ataacacatt	attttgacac	900
aaatgggcag	atctagcaat	ttaaactctg	agtccttcaa	gactgctgtt	cttacgaagt	960
tcacgtccct	gaatcatggt	cctgtatgga	agcctgaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaagggtttac	aaaatatacc	ctgcgggctt	gacacagagg	caagctcttt	1080
atacccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttggagagc	aatccttgtg	1140
ccaagtttga	agtaattttt	gtgtagcata	tgttgagcta	cctacaattt	acatgatcac	1200
ctagcattag	ctcttttact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtaa	tgccctaccct	actttggcca	actcatcggg	gattttacatt	1320
cagaaaatat	acatgacttc	aaccataact	aaaccctttt	ttgtaagata	actgaatggt	1380
catattttaat	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttag	1440
acaacaagat	tgtgggtctg	tactgttatt	tattttattt	tttttttagca	gaaacacctt	1500
atcttttgtt	tcgtttgaat	gtagaatgaa	aataaaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagtttaga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atttattgtg	tttttctttt	1680
ttctaccgga	actttttaga	acggatccca	actcgttccg	gggcccgtac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	accatttcac	1800
catttgattc	agatgtggtg	actagagata	aagcatacta	atttgactct	tggaaaccca	1860
taaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920
tgtgggttgg	tttccattct	gatttatgcg	gcgacttgta	atttaaaatc	taggaggggc	1980
agacattgaa	caatcccaat	atttttaata	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagtg	tcactttgat	agttcaattt	2160
catcttggct	tcttattcct	tttataattc	taattcttct	tgtgtaaaact	atttcattgta	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcctc	actaaactca	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacaa	aaatttacc	aaatttcgac	atcgtttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaatttaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtgtgta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaaact	agttacccta	caacttaaa	aacttaaga	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	taatgttgac	tttgacaccg	2580
caccacatgt	gccttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgtcttatt	2640
ataagcgac	ccctagctcc	aaccaaagga	cgatggagtt	aagaaagaaa	ctcttgetta	2700
cttgtaaggt	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcatctc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttctca	ccccaccatc	catctctcgt	ccc		2863

&lt;210&gt; 6

&lt;211&gt; 2124

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 6

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaaat	aagtcaatct	60
aaacagaccc	taaattccatt	atggtagcta	tcatttttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tattttgaca	caaattgggca	gatctagcaa	180
tttaactctg	gagtccttca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240



tccctgtatgg	aagcctgaaa	gacctcaaat	tctaaaaggt	ggcgataaat	tgaaggttta	300
caaaatatac	cctgcgggct	tgacacagag	gcaagctctt	tataccttcc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttggagag	caatccttgt	gccaagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttactgaga	gaatgaagtt	ttaggaatga	gtatgacct	ggagtcggca	tggtcttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggatttacat	tcagaaaata	tacatgacct	600
caaccatact	taaacccctt	tttgtaagat	aactgaatgt	tcataattta	tggtgggttg	660
tagtggtttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttatttattt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagttagg	atcctctccg	gtcaccggaa	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actattattt	tattttattgt	gtttttcttt	tttctaccgg	aacttttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgtaagtttt	cctttacaga	taccatttca	ccatttgatt	cagatgtggg	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaccc	ataaagttta	tggtatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtgggttg	gtttccattc	1200
tgatttatgc	ggcgacttgt	aattttaaat	ctaggagggg	cagacattga	acaatcccaa	1260
tatttttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	gtgactgaga	1320
ttgagtcata	catttcacta	agaaatgggt	ccaagtacca	aactatcatg	accagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcactctggc	ttcttattcc	1440
ttttataatt	ctaattcttc	ttgtgtaaac	tatttcatgt	attatttttc	tttaaaattt	1500
acatgtcatt	tattttgcct	cactaactca	attttgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaatttcat	caaatttcga	catcgtttat	tatgttcatt	ggatgattaa	1620
caaataatac	aaactttgca	actaattaac	caccaactga	atataattaa	ctataactgt	1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaaatga	attattcata	1740
tatttgaaact	aagttaccct	acaacttaaa	gaacttaaa	aactcgggtt	gagacctggg	1800
gacgaaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgctttttac	1860
atatagttta	tatgacaagt	aatgacaatc	cttgctctat	tataaggcga	cccttagctc	1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgtaagg	tccacacttc	1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac	2040
aaactcattc	caaacacact	ctcttctcca	aaaatgtctt	ccttctcagt	acgtttcctc	2100
acccaccat	ccatctctcg	tccc				2124

&lt;210&gt; 7

&lt;211&gt; 1160

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 7

gatcccaact	cgttccgggg	ccgctacaac	tgaacaaaaa	gaagatattt	tctctctctt	60
cagaaatgta	agttttcctt	tacagatacc	cattcaccat	ttgattcaga	tggtgtgact	120
agagataaag	catactaatt	tgactcttgg	aaaccataa	agtttatgtt	atccgtgttc	180
tggaaccaatc	cacttggggg	cataacctgt	gtctatgtgt	ggtttggttt	ccattctgat	240
ttatgcggcg	acttgtaatt	taaaatctag	gaggggcaga	cattgaacaa	tcccaatatt	300
ttaataactt	atgcaagatt	tttttatta	atgagatgat	gtgtttgtga	ctgagattga	360
gtcatacatt	tcactaagaa	atggttccaa	gtaccaaact	atcatgacct	agttgcaaac	420
atgacgttcg	ggagtgggtca	ctttgatagt	tcaatttcat	cttggcttct	tattcctttt	480
ataattctaa	ttcttcttgt	gtaaaactatt	tcattgtatta	tttttcttta	aaattttacat	540
gtcatttatt	ttgcctcact	aactcaattt	tgcatataac	aatgataagt	gatattttga	600
ctcacaaaat	ttacatcaaa	tttcgacatc	gtttattatg	ttcattggat	gattaacaaa	660
tataacaaa	tttgcaacta	attaaccacc	aactgaatat	aattaactat	aactgtgaaa	720
gtagttaacc	atatttttta	gatgtatata	tcattcgttg	aatgtaatta	ttcatatatt	780
tgaactaagt	taccctacaa	cttaagaact	cggtttgaga	cctggggacg		840
aaaatgtaat	gagactttta	tggtgacttt	gacaccgcac	cacatgtgcc	ttttacatat	900
agtttatatg	acaagtaatg	acaatccttg	ctctattata	aggcgacct	tagtctccac	960
caaaggacga	tggaagttaag	aaagaaactc	ttgcttactt	gtaagggtcca	cactttctca	1020

ctcacctctc aatttcatcc tacaaaaatg tccaaacttc tctttctcac aatcacaaac	1080
tcattccaaa cacactctct tctccaaaaa tgtcttcctt ctcagtagct ttcctcaccc	1140
caccatccat ctctcgtccc	1160

&lt;210&gt; 8

&lt;211&gt; 2904

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 8

ctgtacattc atcttgccgc ctttgcattc acttggccac aaagagtaga gagaaggaag	60
agaagagccc agacttcaag aagcgacctt gcaagtgcac tcgagggtca gaaactgtat	120
atcatatcta tgtgagagaa aggggaacat ttgagatgga gtccatttac ttgaggatata	180
cttattatct tgatcaataa atttgtatac ttcttattta gatcaataaa tttgtcatta	240
agctataatc caaaataaat tacgatcaaa tatgcaaatg ttagccagta cttgtgttaa	300
acttgatggc atctcttggg ttctttggca atcacatgcc taagaaataa atagtatcat	360
atgatttgtt ttggtcagac ttcagagtca gatgactctg tttggataaa cagcttaatt	420
aagcgcttat agaatatcat atgatttgtt ttggtcagac ttcagagcat ctcttgggtt	480
ctctggcaat catatgccta agaaataaat agtatcatat gatttgtgtt ggtcagactt	540
cagagtcaga tgacctgtt tgggtaaaaca gcttaattaa gtgcttatag aataagcgct	600
tatcatataa gtgcttttgt acagttattt ctatgaaagt agaagaaata gtcatattgt	660
tttaatatata gctatcctgg agagcttggg gaaataacca gaaaagaact tatggacacg	720
tcagtagctg tttacataag atctccctaa cagtctcaaa agtgcttatg ccagtagata	780
aattcaataa agtcaatcta aacagaccct aaatccatta tggtagctat catttttagct	840
tattccatct ttattaagaa tgtcatgaga taacataatg ataacacatt attttgacac	900
aaatgggcag atctagcaat ttaactctgg agtccctcaa gactgctgtt cttacgaagt	960
tcacgtccct gaatcatgtt cctgtatgga agcctgaaag acctcaaat ctaaaagggtg	1020
gcgataaatt gaaggtttac aaaatatacc ctgcgggctt gacacagagg caagctcttt	1080
ataccttcca gttcaacggg gatgttgatt tcagaagtca cttggagagc aatccttggtg	1140
ccaagtttga agtaattttt gtgtagcata tgttgagcta cctacaattt acatgatcac	1200
ctagcattag ctctttcact taactgagag aatgaagttt taggaatgag tatgaccatg	1260
gagtgcggca gcctttgtaa tgctaccct actttggcca actcatcggg gatttacatt	1320
cagaaaatat acatgacttc aaccataact aaaccccttt ttgtaagata actgaatgtt	1380
catatttaat gttgggttgt agtggtttta cttgattata tccagacagt tacaagttgg	1440
acaacaagat tgtgggtctg tactgttatt tatttatttt ttttttagca gaaacacctt	1500
atcttttgtt tcgtttgaat gtagaatgaa aataaaaagaa agaaaatata acatcatcgg	1560
ccgcgcttgt ctaatttcgg gcagtttaga tcctctccgg tcaccggaaa gtttcagtag	1620
aagaaacaaa acacgtgac taaaatgata ctattatttt atttatttgt tttttctttt	1680
ttctaccgga acttttttaga acggatccca actcgttccg gggccgctac aactgaaaca	1740
aaagaagata ttttctctct cttcagaaat gtaagttttc ctttacagat acccattcac	1800
catttgattc agatgtggtg actagagata aagcatacta atttgactct tggaaacca	1860
taaaagttat gttatccgtg ttctggacca atccacttgg gggcataacc tgtgtctatg	1920
tgtggttttg tttccattct gatttatgcg gcgacttgta atttaaaatc taggaggggc	1980
agacattgaa caatcccaat attttaataa cttatgcaag atttttttta ttaatgagat	2040
gatgtgtttg tgactgagat tgagtcatac atttcactaa gaaatgggtc caagtaccaa	2100
actatcatga ccagttgca aacatgacgt tcgggagtggt tcactttgat agttcaattt	2160
catcttggct tcttattcct tttataattc taattcttct tgtgtaaaact atttcatgta	2220
ttatttttct ttaaaattta catgtcattt attttgctc actaactcaa ttttgcatat	2280
aacaatgata agtgatattt tgactcacaa aatttacatc aaatttcgac atcgtttatt	2340
atgttcattg gatgattaac aaatataaca aactttgcaa ctaattaacc accaactgaa	2400
tataattaac tataactgtg aaagtagtta accatatttt ttagatgtat atatcatccg	2460
ttgaatgtaa ttattcatat atttgaacta agttacccta caacttaaag aacttaaaga	2520
actcggtttg agacctgggg acgaaaatgt attgagactt caatgttgac tttgacaccg	2580
caccacatgt gccttttaca tatagtttat atgacaagta atgacaatcc ttgctctatt	2640
ataaggcgac ccttagctcc aaccaaagga cgatggagtt aagaaagaaa ctcttgctta	2700
cttgtaaggt ccacacttct tcactcacct ctcaatttca tcctacaaa atgtccaaac	2760

ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttcctca	ccccaccatc	catctctcgt	cccaacaaaa	catggctact	2880
atctgctgca	actccatcag	ttgc				2904

&lt;210&gt; 9

&lt;211&gt; 2165

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 9

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaaat	aagtcaatct	60
aaacagacc	taaatccatt	atggtagcta	tcatttttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tattttgaca	caaatgggca	gatctagcaa	180
tttaactctg	gagtccttca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240
tcctgtatgg	aagcctgaaa	gacctcaaat	tctaaaaggt	ggcgataaat	tgaagggttta	300
caaaatatac	cctgcgggct	tgacacagag	gcaagctctt	tataccttcc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttggagag	caatccttgt	gccaagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttaaactgaga	gaatgaagtt	ttaggaatga	gtatgaccat	ggagtgggca	tggctttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggattttacat	tcagaaaata	tacatgactt	600
caaccatact	taaaacctt	tttgaagat	aactgaatgt	tcataatttaa	tgttggggtg	660
tagtggtttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttattttatt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagttagg	atcctctccg	gtcaccggaa	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actattattt	tattttattgt	gtttttctt	tttctaccgg	aactttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgtgaagttt	cctttacaga	taccatttca	ccatttgatt	cagatgtggt	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaccc	ataaagttta	tgttatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtgggttg	gtttccattc	1200
tgattttatgc	ggcgacttgt	aattttaa	ctaggagggg	cagacattga	acaatcccaa	1260
tatttttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	gtgactgaga	1320
ttgagtcata	catttcacta	agaaatggtt	ccaagtacca	aactatcatg	accagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcactctggc	ttcttattcc	1440
ttttataatt	ctaattcttc	ttgtgtaa	tatttcattg	attatttttc	tttaaaattt	1500
acatgtcatt	tattttgcct	cactaactca	attttgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaattttacat	caaatttcga	catcgtttat	tatgttcatt	ggatgattaa	1620
caaatataac	aaactttgca	actaattaac	caccaactga	atataattaa	ctataactgt	1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaatgta	attattcata	1740
tatttgaact	aagttaccct	acaacttaaa	gaacttaaa	aactcgggtt	gagacctggg	1800
gacgaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgccttttac	1860
atatagttta	tatgacaagt	aatgacaatc	cttgctctat	tataaggcga	cccttagctc	1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgtaagg	tccacacttc	1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac	2040
aaactcattc	caaacacact	ctcttctcca	aaaatgtctt	ccttctcagt	acgtttcctc	2100
acccacccat	ccatctctcg	tcccaacaaa	acatggctac	tatctgctgc	aactccatca	2160
gttgc						2165

&lt;210&gt; 10

&lt;211&gt; 1316

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 10

gatcctctcc	ggtcaccgga	aagtttcagt	agaagaaaca	aaacaccgtg	actaaaatga	60
tactattatt	ttattttattg	tgtttttctt	ttttctaccg	gaacttttta	gaacggatcc	120
caactcggtc	cggggccgct	acaactgaaa	caaaagaaga	tattttctct	ctcttcagaa	180
atgtaagttt	tcctttacag	ataccattc	accatttgat	tcagatgtgg	tgactagaga	240
taaagcatac	taatttgact	cttggaacc	cataaagttt	atgttatccg	tggtctggac	300
caatccactt	gggggcataa	cctgtgtcta	tggtgtggtt	ggtttccatt	ctgatttatg	360
cggcgacttg	taatttaaaa	tctaggaggg	gcagacattg	aacaatccca	atattttaat	420
aacttatgca	agattttttt	tattaatgag	atgatgtgtt	tgtgactgag	attgagtcac	480
acatttcact	aagaaaatgg	tccaagtacc	aaactatcat	gaccagttg	caaactgac	540
gttcgggagt	ggtcactttg	atagttcaat	ttcatcttgg	cttcttatcc	cttttataat	600
tctaattctt	cttggtgtaa	ctatttcatg	tattattttt	ctttaaaatt	tacatgtcat	660
ttattttgcc	tacttaactc	aattttgcat	ataacaatga	taagtatat	tttgactcac	720
aaaattttaca	tcaaatttcg	acatcgttta	ttatgttcat	tggatgatta	acaaatataa	780
caaaactttgc	aactaattaa	ccaccaactg	aataataatta	actataactg	tgaagtagt	840
taaccatatt	ttttagatgt	atataatcat	cgttgaatgt	aattattcat	atatttgaac	900
taagttaccc	tacaacttaa	agaacttaaa	gaactcggtt	tgagacctgg	ggacgaaaa	960
gtaatgagac	tttaatgttg	actttgacac	cgcaccacat	gtgcctttta	catatagttt	1020
atatgacaag	taatgacaat	ccttgctcta	ttataaggcg	acccttagct	ccaaccaaag	1080
gacgatggag	ttaagaaaga	aactcttgct	tacttgtaag	gtccacactt	cttactcac	1140
ctctcaattt	catcctacaa	aaatgtccaa	acttctcttt	ctcacaatca	caaaactcatt	1200
ccaaacacac	tctcttctcc	aaaaatgtct	tccttctcag	tacgtttcct	cacccaccca	1260
tccatctctc	gtcccaacaa	aacatggcta	ctatctgctg	caactccatc	agttgc	1316

&lt;210&gt; 11

&lt;211&gt; 2971

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 11

ctgtacattc	atcttgccgc	ctttgcattc	acttgggccac	aaagagtaga	gagaaggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgaggggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccatttac	ttgagggtata	180
cttatttttt	tgatcaataa	atttgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaag	ttagccagta	cttggtgtaa	300
acttgatggc	atctcttggt	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgatttgtt	ttgggtcagac	ttcagagtca	gatgactctg	tttggtataa	cagcttaatt	420
aagcgcttat	agaatatcat	atgatttgtt	ttgggtcagac	ttcagagcat	ctcttggttt	480
ctctggcaat	catatgecta	agaaataaat	agtatcatat	gattgtgttt	ggtcagactt	540
cagagtcaga	tgacctgtt	tgggtaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttattt	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaataata	gctatcctgg	agagcttggt	gaaataacca	gaaaagaact	tatggacacg	720
tcagagctg	tttacataag	atctccctaa	cagtctcaaa	agtgtttatg	ccagtagata	780
aattcaaata	agtcaatcta	aacagacctt	aaatccatta	tggtacctat	catttttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taacataatg	ataacacatt	attttgacac	900
aatgggag	atctagcaat	ttaactctgg	agtccttcaa	gactgctgtt	cttacgaagt	960
tcacgtccct	gaatcatgtt	cctgtatgga	agcctgaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaagggtttac	aaaatatacc	ctgctgggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttgagagac	aatccttggtg	1140
ccaagtttga	agtaattttt	gtgtagcata	tggttagagta	cctacaattt	acatgatcac	1200
ctagcattag	ctctttcact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtta	tgccctaccct	actttggcca	actcatcggtg	gatttacatt	1320
cagaaaatat	acatgacttc	aaccatactt	aaaccccttt	ttgtaagata	actgaatgtt	1380
catattttaat	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttgg	1440
acaacaagat	tgtgggtctg	tactgttatt	tattttattt	tttttttagca	gaaacacctt	1500

atcttttgtt	tcgtttgaat	gtagaatgaa	aataaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagtttaga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atttattgtg	tttttctttt	1680
ttctaccgga	acttttttaga	acggatccca	actcgttccg	gggcccgtac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	acccattcac	1800
catttgattc	agatgtggtg	actagagata	aagcactacta	atttgactct	tggaaaccca	1860
taaaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920
tgtggttttg	tttccattct	gattttatgcg	gcgacttgta	atttaaaatc	taggaggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	ccaggttgca	aacatgacgt	tcgggagtg	tcactttgat	agttcaattt	2160
catcttggtc	tcttattcct	tttataattc	taattcttct	tgtgtaaact	atttcatgta	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcctc	actaactcaa	ttttgcata	2280
aacaatgata	agtgatattt	tgactcacaa	aatttacatc	aaatttcgac	atcgtttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaacta	agttacccta	caacttaaa	aacttaaa	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	taatggtgac	tttgacaccg	2580
caccacatgt	gccttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgctctatt	2640
ataaggcgac	ccttagctcc	aaccaaagga	cgatggagtt	aagaaagaaa	ctcttgctta	2700
cttgaaggt	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttctcga	cccaccatc	catctctcgt	cccaacaaaa	catggctact	2880
atctgctgca	actccatcag	ttgcacctgt	ttcaacacca	caagttgatg	catcaaggtt	2940
ggagcctaga	gttgaggaaa	aagatggtta	c			2971

&lt;210&gt; 12

&lt;211&gt; 2232

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 12

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaaat	aagtcaatct	60
aaacagaccc	taaatccatt	atggtagcta	tcatttttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tattttgaca	caaattgggca	gatctagcaa	180
tttaactctg	gagtcttcca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240
tcctgtatgg	aagcctgaaa	gacctcaaat	tctaaaaggt	ggcgataaat	tgaagggtta	300
caaaatatac	cctgcccggc	tgacacagag	gcaagctctt	tataccttcc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttggagag	caatccttgt	gccaagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttaactgaga	gaatgaagtt	ttaggaatga	gtatgaccat	ggagtcggca	tggctttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggattttacat	tcagaaaata	tacatgactt	600
caaccatact	taaacccctt	tttgtaagat	aactgaatgt	tcataatttaa	tgttggggtg	660
tagtgttttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttattttatt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagttagg	atcctctccg	gtcacccgga	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actattattt	tattttattgt	gtttttcttt	tttctaccgg	aacttttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgaagtttt	cctttacaga	taccatttca	ccatttgatt	cagatgtggg	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaccc	ataaagttta	tgttatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtgggttg	gtttccattc	1200
tgattttatgc	ggcgacttgt	aattttaaaat	ctaggagggg	cagacattga	acaatcccaa	1260
tatttttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	gtgactgaga	1320
ttgagtcata	catttcacta	agaaatggtt	ccaagtacca	aactatcatg	acccagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcactctggc	ttcttattcc	1440

ttttataatt	ctaattcttc	ttgtgtaaac	tatttcatgt	attatttttc	tttaaaat	1500
acatgtcatt	tattttgcct	cactaactca	attttgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaattttacat	caaatttcga	catcgtttat	tatgttcatt	ggatgattaa	1620
caaataatac	aaactttgca	actaatga	caccaactga	atataattaa	ctataactgt	1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaatgta	attattcata	1740
tatttgaact	aagttaccct	acaacttaaa	gaacttaaag	aactcgggtt	gagacctggg	1800
gacgaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgctttttac	1860
atatagttta	tatgacaagt	aatgacaatc	cttgctctat	tataaggcga	cccttagctc	1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgtaagg	tccacacttc	1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac	2040
aaactcattc	caaacacact	ctcttctcca	aaaatgtctt	ccttctcagt	acgtttcttc	2100
acccaccat	ccatctctcg	tcccaacaaa	acatggctac	tatctgctgc	aactccatca	2160
gttgacactg	tttcaacacc	acaagttgat	gcacaaagg	tggagcctag	agttgaggaa	2220
aaagatgggt	ac					2232

&lt;210&gt; 13

&lt;211&gt; 1383

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as promoter for regulating expression of foreign genes

&lt;400&gt; 13

gacacctctc	ggtcaccgga	aagtttcagt	agaagaaaca	aaacaccgtg	actaaaatga	60
tactattatt	ttatttattg	tgtttttctt	ttttctaccg	gaacttttta	gaacggatcc	120
caactcggtc	cggggccgct	acaactgaaa	caaaagaaga	tattttctct	ctcttcagaa	180
atgtaagttt	tcctttacag	ataccattc	accatttgat	tcagatgtgg	tgactagaga	240
taaagcatac	taatttgact	cttggaacc	cataaagttt	atgttatccg	tgttctggac	300
caatccactt	gggggcataa	cctgtgtcta	tggtgggttt	ggtttcatt	ctgatttatg	360
cggcgacttg	taatttaaaa	tctaggagg	gcagacattg	aacaatccca	atattttaat	420
aacttatgca	agattttttt	tattaatgag	atgatgtgtt	tgtagctgag	attgagtcac	480
acatttcact	aagaaatggg	tccaagtacc	aaactatcat	gaccagttg	caaacatgac	540
gttcgggagt	ggtcactttg	atagttcaat	ttcatcttgg	cttcttatcc	cttttataat	600
tctaattctt	cttgtgtaaa	ctatttcatt	tattattttt	ctttaaaatt	tacatgtcat	660
ttattttgcc	tcactaactc	aattttgcat	ataacaatga	taagtgatat	tttgactcac	720
aaaatttaca	tcaaatttcg	acatcgttta	ttatgttcat	tgtagtatta	acaatatata	780
caaactttgc	aactaattaa	ccaccaactg	aataataatta	actataactg	tgaaagtagt	840
taaccatatt	ttttagatgt	atataatcat	cgttgaaatg	aattattcat	atatttgaac	900
taagttaccc	tacaacttaa	agaacttaaa	gaactcgggt	tgagacctgg	ggacgaaaat	960
gtaatgagac	tttaatgttg	actttgacac	cgcaccacat	gtgcctttta	catatagttt	1020
atatgacaag	taatgacaat	ccttgctcta	ttataaggcg	acccttagct	ccaaccaaaag	1080
gacgatggag	ttaagaaaga	aactcttgct	tacttgtaag	gtccacactt	cttcactcac	1140
ctctcaattt	catcctacaa	aaatgtccaa	acttctcttt	ctcacaatca	caaactcatt	1200
ccaaacacac	tctcttctcc	aaaaatgtct	tccttctcag	tacgtttcct	cacccaccca	1260
tccatctctc	gtcccaacaa	aacatgggta	ctatctgctg	caactccatc	agttgcacct	1320
gtttcaacac	cacaagttga	tgcatcaagg	ttggagccta	gagttgagga	aaaagatggg	1380
tac						1383

&lt;210&gt; 14

&lt;211&gt; 3472

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as terminator for regulating expression of foreign genes

&lt;400&gt; 14

caagttgatg	catcaagggt	ggagcctaga	gttgaggaaa	aagatgggta	ctggggtttg	60
aaggaagagt	atagaggagg	tattaatcct	caggagaaag	ttaagattca	gaaagaacct	120
atgaagcttt	ttatggaagg	tgggattaat	gatttggcta	atatgtctct	tgaagagatt	180
gaaagctcta	agcttactaa	agatgatatt	gatgttagac	ttaaatggct	tggtcttttt	240
catagaagga	aacatcattg	taagtttttt	tactttcttt	ttatacttca	aagttctctc	300
atactctgta	tttggtttatt	agttttttgta	gacttaaata	ttctctttga	tttacctagt	360
gaaactccat	tcttggtttcc	gaaattgtag	tgtgtatagt	ctagaaaatt	aagaagtaga	420
caaaatgatt	tatgagattg	taaattgtag	gctttttatc	aattttattaa	ttttagagac	480
caaaatttgc	ctatcttatt	tggaccaatt	tattgtatgt	taggatcgac	atgagtttag	540
caaaatcacc	acggcaccat	gactgtgttg	aagcttcttt	gtgtaacttt	aacaaaaatt	600
atatggcaca	ccatgattat	gcaaactcac	cgtcaatcca	aacatagaaa	ttcagtgtta	660
atctttgtga	caataaaaaa	ctatgagtta	tgttgtacta	atttatttcc	attgtgaaac	720
tcagatggta	gatttatgat	gagactaaaa	ctcccaaatg	gggtaacaac	aagtgtctca	780
acaagatact	tggcgagtgt	gataaaaaaa	tatggcaaa	acggatgtgc	tgatgtgaca	840
acgaggcaga	attggcaaat	tcgaggtgta	acgttacctg	atgtccctga	aattcttaag	900
ggccttgcag	aggtcggctt	gacaagctctg	cagagtggaa	tggacaatgt	tcgaaacca	960
gttggttaacc	ctcttgctgg	tattgacctt	gatgagattg	ttgatacaag	accttacacc	1020
aatttgctgt	cccaattcat	cactgcta	tcacttggta	atccaacccat	tacaaacttg	1080
taagtctaaa	ctatctcatc	tttatatttc	actcattata	tcataattagt	agtttagttac	1140
ttgcattgca	agcattacgt	gaccgtgtgt	agcctctaaa	tccttttgat	aatatgtgca	1200
ggccaaggaa	gtggaatgta	tgtgtgatag	gttcccatga	tcttttcgag	catccgcata	1260
ttaacgatct	tgtttatatg	cctgctaata	aggatggctg	atttggattc	aacttattgg	1320
tgggtgggtt	cttttagtccc	aagcgatgtg	ctgaagcagt	tccacttgat	gcatgggtct	1380
ctgcagatga	tgttatccca	ctttgtaaa	ctgtccttga	gacctatagg	gacctcgga	1440
caagagggaa	tagacagaaa	accagaatga	tgtgggtgat	cgatgaactt	gtaagttacc	1500
actttttttc	ttcacatatt	attaactgaa	gtgacttta	cgaccatttt	acaattgaaa	1560
tttaagtggg	ttttagccct	atcattacaa	gaacaaattt	gttaattcac	tagcaagagc	1620
aattccactt	tggcttggac	atgacaagt	tttgtgaaat	gcaggggata	gaagtattca	1680
gatcagaggt	ggaaaaaaga	atgccagaga	agaagctaga	gagagcatcc	aaagaagaac	1740
ttgtccaaaa	acaatggaaa	gaggagacat	cttaggtgtt	catccacaaa	aacaagaagg	1800
tttaagctat	gttggaattc	acattccagt	tggtagaatc	caagcagatg	agatggaaga	1860
gctagctcgt	atcgccgatg	aatacggaa	cggagaacta	aggctaaccg	tggagcaaaa	1920
cataataatt	ccaaatgtgg	aaaactcaaa	acttgatgca	ttgctaata	aacctctctt	1980
gaaagacaaa	ttctcaccag	aaccttccat	cctaataaaa	acacttgtgg	catgcactgg	2040
taaccaattt	tgtggccaag	caataattga	aacaaaaaaa	agagctttta	aagtaactga	2100
agaagtggaa	agacatgtgg	ctgtgagcaa	accagtgaga	atgcattgga	ctgggtgtcc	2160
taacacttgt	ggtcaagtcc	aggttgctga	tattgggttt	atgggttgta	tggctaggga	2220
tgagaatggt	aaggctactg	aagggtgtga	tattttcctt	gggtgggagaa	ttggaagtga	2280
ttctcattta	gctgaggtgt	ataagaaagg	tgtcccttgc	aaggacttgg	tgcctattgt	2340
agctgatatt	ttggttaaat	attttggagc	tgtccaaaag	aatagagaag	aaggggatga	2400
ttaaagtata	taggtatttg	gtgattttta	ttgcctctac	acaaaattat	tatgttctgt	2460
ccaaaatata	aagtcacaag	ggataattga	gattgagatg	cagcacgcca	cacatgaact	2520
tgtacatttg	gataagtcac	ttttcattgc	tattttataa	gttacacttt	gaattttata	2580
ataaatttta	ttttatttca	aggaccagat	tttataagga	aaccgcta	ctaactatct	2640
ttactcgtaa	tttgtcattt	gagagctacg	gagatcggtg	agtttacgta	tgagtgttta	2700
gtctcacatt	aattatgaat	ggtcaaaatg	ttaaatttat	aagagatgta	atctatatac	2760
ctaattgcatt	aaaaatttgg	atggagatgc	gacgcccccc	ttttttgtgg	tcctgaagta	2820
tagacttggt	gtcgcttctg	gtgcactctc	atacttccca	acaaggagaa	aaaactacca	2880
taacaattaa	caaactaaca	tttggtattt	aaaaaaacat	acggatactg	ttttttcccc	2940
atttatttag	aagatgatgg	cttggatttc	aatggctgag	tttatttttt	ttttggtegg	3000
gagttgaagt	atcgggaaaa	ctaaaatgct	tatgacttta	aacattgtgt	tgatatatga	3060
ttagttttca	acttacttaa	aaagtggcaa	actagtttag	tggttctctc	ccttccttgt	3120
agttcaagga	acatgggttt	gaactctgtc	caaatttttg	tactttcaat	tatccataca	3180
tttaaaagct	atataccaca	tcattatatt	caagtcaatg	atcatgcggc	ctgccacatt	3240
agcatcgatg	tacacattaa	ttttaagtgg	catgaacaca	ttaacatttc	ataaaagcta	3300
tgtgccagat	catcattcaa	gtctatgcac	acatgggtcaa	cacattagta	ccattttttt	3360
ttttattggt	gatcagattg	gatgtcggta	ttgtgtgtat	gtacaaaact	caaacaatct	3420
ccagctgtta	gagaacgtcg	aaaatgaatg	catcacgggt	gcacacttag	at	3472

&lt;210&gt; 15

13/17

<211> 617  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Sequences to be used as terminator for regulating  
 expression of foreign genes

<400> 15

agtatatagg	tatttggtga	ttttaattgc	ctctacacaa	aattattatg	ttctgtccaa	60
aatataaagt	cacaagggat	aattgagatt	gagatgcagc	acgccacaca	tgaacttgta	120
catttggata	agtcattttt	cattgctatt	ttataagtta	cactttgaat	tttataataa	180
attttatttt	atttcaagga	ccagatttta	taaggaaaacc	gctaattctaa	ctatctttac	240
tcgtaatttg	tcatttgaga	gctacggaga	tcgttgagtt	tacgtatgag	tgtttagtct	300
cacattaatt	atgaatggtc	aaaatgttaa	atttataaga	gatgtaatct	atatacctaa	360
tgcattaaaa	atttggatgg	agatgcgacg	ccccctttt	ttgtggtcct	gaagtataga	420
cttggtgtcg	cttctggtgc	actctcatat	ttcccaacaa	ggagaaaaaa	ctaccataac	480
aattaacaaa	ctaacatttg	ttatttaaaa	aaacatacgg	atactgtttt	ttccccattt	540
attaggaaga	tgatggcttg	gatttcaatg	gctgagttta	tttttttttt	ggtcgggag	600
tgaagtatcg	ggaaaac					617

<210> 16  
 <211> 502  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Sequences to be used as terminator for regulating  
 expression of foreign genes

<400> 16

ttgtacattt	ggataagtca	tttttcattg	ctattttata	agttacactt	tgaattttat	60
aataaatttt	attttatttc	aaggaccaga	ttttataagg	aaaccgctaa	tctaactatc	120
tttactcgta	atttgtcatt	tgagagctac	ggagatcggt	gagtttacgt	atgagtgttt	180
agtctcacat	taattatgaa	tggtcaaaat	gttaaattta	taagagatgt	aatctatata	240
cctaattgcat	taaaaatttg	gatggagatg	cgacgcccc	cttttttggtg	gtcctgaagt	300
atagacttgt	tgtcgcttct	ggtgcactct	catacttccc	aacaaggaga	aaaaactacc	360
ataacaatta	acaaactaac	atttgttatt	taaaaaaaca	tacggatact	gttttttccc	420
catttattag	gaagatgatg	gcttggattt	caatggctga	gtttattttt	tttttggtcg	480
ggagttgaag	tatcgggaaa	ac				502

<210> 17  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Sequences to be used as primers

<400> 17

gatattgatg ttagactcaa gtggc

25

<210> 18  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence



&lt;220&gt;

&lt;223&gt; Sequences to be used as primers

&lt;400&gt; 18

cacysattcc acttcctwgg c

21

&lt;210&gt; 19

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as primers

&lt;400&gt; 19

ttgtcacatc agcacatccg tcttttgc

27

&lt;210&gt; 20

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as primers

&lt;400&gt; 20

tcgccaaagta tcttggtttga gcacttg

27

&lt;210&gt; 21

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as primers

&lt;400&gt; 21

agagcccggg agaagagagt gtgtttg

27

&lt;210&gt; 22

&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Sequences to be used as primers

&lt;400&gt; 22

ttctcccggg ggacgagaga tggatggt

28

<210> 23  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sequences to be used as primers

<400> 23  
ttctcccggg gttgaaacag gtgcaactga 30

<210> 24  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sequences to be used as primers

<400> 24  
ttctcccggg taaccatctt tttcttca 28

<210> 25  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sequences to be used as primers

<400> 25  
cacactttctt cactcacctc tcaa 24

<210> 26  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sequences to be used as primers

<400> 26  
atctaggagg ggcagacatt g 21

<210> 27  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> sequences to be used as primers

<400> 27

tcggtataaa gacttcgcgc tgat 24

<210> 28  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> sequences to be used as primers

<400> 28  
atgtcttcct tctcagtagc tttcctc 27

<210> 29  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> sequences to be used as primers

<400> 29  
caagttgatg catcaaggtag ggagcctaga 30

<210> 30  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> sequences to be used as primers

<400> 30  
agaagagctc agtatatagg tatttggtga 30

<210> 31  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> sequences to be used as primers

<400> 31  
agaagagctc ttgtacattt ggataagtca 30

<210> 32  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> sequences to be used as primers

<400> 32  
agaagaattc gttttcccga tacttcaact

30

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**